

JUMBOTM

Positioning chair

JUMBOTM chair is ideal for children with both increased muscle tone and very flaccid muscle tone. It supports daily activities like rehabilitation exercises, eating, and playing. Its MEDICUS™ medical upholstery is vapor-permeable, waterproof, and easy to disinfect, making it suitable for home use, hospitals, or rehabilitation centers. The foldable footrest aids in seating and can be easily removed for storage and transport, with adjustable length settings for flexibility. Made from eco-friendly, safe materials, the JUMBOTM chair is especially recommended for individuals with difficulty maintaining a sitting position, particularly those with neurological or orthopedic conditions.



Lightweight

24⁺

24-month
warranty



Stable
construction



Easy
operation

Colors

MEDICUS™ upholstery


Waterproof and steam-permeable



Med Sea



Independently adjustable side supports ensure proper trunk stabilization.



Due to available adjustments we are able to set the desired seat depth and tilt angle of the backrest.



110 - 180 cm

Indicative user
height



30 - 75 kg

Max. user
weight



For **more details** about the product, please visit our website.

This is a medical device.
Use it in accordance with the user manual or the label.

INDICATIONS

Cerebral palsy (CP) – various types and forms | Brain and spinal cord injuries of various etiologies with limb paresis, severe balance disturbances, and coordination impairments | Demyelinating diseases (multiple sclerosis – MS, amyotrophic lateral sclerosis – ALS, and others) | Meningocele | Muscle diseases with paresis and paralysis (including dystrophy, myasthenia gravis, polymyositis) | Genetic syndromes with limb paresis | Degenerative and metabolic diseases of the nervous system | Degenerative and rheumatic joint diseases | Other diseases with limb paresis, paralysis, and musculoskeletal impairments | After injuries, following traumatic brain injuries or spinal cord injuries with limb paresis and paralysis, particularly with quadriplegia (as part of the rehabilitation process or as permanent orthopedic support)

CONTRAINDICATIONS

Observation: Monitor user during standing. Stop therapy if paleness, sweating, fainting, joint overload, or body deformations occur; consult with a doctor or physiotherapist | **Absolute Contraindication:** Critical condition of the user. | **Relative Contraindications:** Resting hypertension > 200/110 mmHg, pulmonary hypertension, heart rhythm disorders, unstable asthma, unstable metabolic issues, moderate heart valve disease. | Conditions like healing fractures or severe osteoporosis that prevent weight-bearing | Incomplete fracture healing (unless approved by a doctor) | Postoperative joint load restrictions | Pain from dislocation, excessive tissue stretch, or pressure; dislocation without pain is not contraindicated | Significant hip/knee flexion or ankle contractures; ensure no excessive pressure or stretch on affected areas.