

# LIFTER™

## Static standing frame

Standing frame LIFTER™ is a modern rehabilitation device. Electric actuator with integrated sit allows the user for independent use of the device by means of the remote control. Powerful actuator can lift the users with weight up to 120 kg. There is a possibility to equip LIFTER™ with upper torso stabilizer which perfectly immobilizes the chest. This solution is highly recommended for users with upper limbs paresis. LIFTER™'s seat also serves as a pelvic support that keeps the user in the proper position and secures him in the device.



Easy  
operation

24<sup>+</sup>

24-month  
warranty



Stable  
construction



Ergonomically shaped and adjustable knee supports perfectly stabilize the lower limbs.



Modern upright standing system uses the lifting mechanism with movable arm, electric actuator and remote control to help in adoption of the upright standing position.




120 - 200 cm

Indicative user  
height



120 kg

Max. user  
weight



For **more details** about  
the product, please visit  
our website.

**This is a medical device.**  
**Use it in accordance with the user manual or the label.**

### INDICATIONS

Cerebral palsy (CP) – various types and forms | Brain and spinal cord injuries of various etiologies with limb paresis, severe balance disturbances, and coordination impairments | Demyelinating diseases (multiple sclerosis – MS, amyotrophic lateral sclerosis – ALS, and others) | Meningocele | Muscle diseases with paresis and paralysis (including dystrophy, myasthenia gravis, polymyositis) | Genetic syndromes with limb paresis | Degenerative and metabolic diseases of the nervous system | Degenerative and rheumatic joint diseases | Other diseases with limb paresis, paralysis, and musculoskeletal impairments | After injuries, following traumatic brain injuries or spinal cord injuries with limb paresis and paralysis, particularly with quadriplegia (as part of the rehabilitation process or as permanent orthopedic support)

### CONTRAINDICATIONS

**Observation:** Monitor user during standing. Stop therapy if paleness, sweating, fainting, joint overload, or body deformations occur; consult with a doctor or physiotherapist | **Absolute Contraindication:** Critical condition of the user. | **Relative Contraindications:** Resting hypertension > 200/110 mmHg, pulmonary hypertension, heart rhythm disorders, unstable asthma, unstable metabolic issues, moderate heart valve disease. | Conditions like healing fractures or severe osteoporosis that prevent weight-bearing | Incomplete fracture healing (unless approved by a doctor) | Postoperative joint load restrictions | Pain from dislocation, excessive tissue stretch, or pressure; dislocation without pain is not contraindicated | Significant hip/knee flexion or ankle contractures; ensure no excessive pressure or stretch on affected areas.